



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

concentrated sulphuric acid, the mineral is decomposed very quickly as soon as the temperature is brought to the boiling point of sulphuric acid. The mass becomes a dry paste and dissolves in water. The solution is turbid from a quantity of basic phosphates, varying between two and eighteen per cent., according to the excess of acid present.

The acid solution may be boiled without the forming of a precipitate; thorium is therefore not contained in the mineral. Two determinations of the phosphoric acid gave 25.82 and 26.3 per cent., one being by phosphomolybdic acid; the other in the usual manner, after precipitating the bases first by oxalic acid, and the filtrate by ammoniac hydrate. Fluorine is not present.

The following is given as a preliminary result, pending the tedious separation of the oxides:

(Ce, La, Dy, Y) ₂ O ₃	=	73.82
(Y, Fe, Ca) ₂ O ₃	=	1.00
P ₂ O ₅	=	26.05
Volatile by ignition	=	0.45
		<hr/>
		101.32

Supposing the oxides to be all cerous oxide, or in other words having the atomic weight of 92, the highest of the group, then the ratio obtains

$$P_2 O_5 : 3 Ce O = 1 : 3.75,$$

which is not reconcilable with a normal phosphate.

The speaker suggests, therefore, the possible presence in the group of a metal with a much higher atomic weight than cerium. He is engaged at work with a large enough quantity of the oxides to decide this question in time.

JANUARY 31.

The President, Dr. LEIDY, in the chair.

Eighteen persons present.

Messrs. Wilson Mitchell, Chas. H. Hutchinson, Rev. W. G. Holland, Able F. Price, Alfred C. Harrison and Robt. B. Haines were elected members.

Dr. A. Baltzer, of Zurich, and Prof. Robt. Collett, of Christiania, were elected correspondents.

The following were ordered to be published:—